


Steven M. Ruben
Appl. No. 10/662,429



BEST AVAILABLE COPY

Ruben EXHIBIT 2108
Ruben v. Wiley et al.
Interference No. 105,077
RX 2108

Molecular Cloning and Expression of the Fas Ligand, a Novel Member of the Tumor Necrosis Factor Family

Takashi Suda,* Tomohiro Takahashi,*
Pierre Golstein,* and Shigekazu Nagata*

*Osaka Bioscience Institute

6-2-4 Furuedai

Suita, Osaka 565

Japan

†Centre d'immunologie Institut National
de la Santé et de la Recherche Médicale

Centre National de la Recherche

Scientifique de Marseille-Luminy

Case 906

13288 Marseille Cedex 9

France

Summary

The Fas antigen (Fas) belongs to the tumor necrosis factor (TNF)/nerve growth factor receptor family, and it mediates apoptosis. Using a soluble form of mouse Fas, prepared by fusion with human immunoglobulin Fc, Fas ligand was detected on the cell surface of a cytotoxic T cell hybridoma, PC60-d10S. A cell population that highly expresses Fas ligand was sorted using a fluorescence-activated cell sorter, and its cDNA was isolated from the sorted cells by expression cloning. The amino acid sequence indicated that Fas ligand is a type II transmembrane protein that belongs to the TNF family. The recombinant Fas ligand expressed in COS cells induced apoptosis in Fas-expressing target cells. Northern hybridization revealed that Fas ligand is expressed in activated splenocytes and thymocytes, consistent with its involvement in T cell-mediated cytotoxicity and in several nonlymphoid tissues, such as testis.

Introduction

Mammalian development is tightly regulated not only by the proliferation and differentiation of cells but also by cell death (Ellis et al., 1991; Raff, 1992). The cell death that occurs during development or tissue turnover is called programmed cell death, most of which proceeds via apoptosis (Walker et al., 1988; Wyllie et al., 1980). Apoptosis is morphologically distinguished from necrosis, which occurs during the accidental cell death caused by physical or chemical agents. During apoptosis, the cytoplasm of the affected cells condenses, and the nucleus also condenses and becomes fragmented. At the final stage of apoptosis, the cells themselves are fragmented (apoptotic bodies) and are phagocytosed by neighboring macrophages and granulocytes (Raff, 1992; Walker et al., 1988; Wyllie et al., 1980). Apoptosis occurs not only during programmed cell death, but also during the death process induced by some cytotoxic T cells, such as tumor necrosis factor (TNF) or lymphotoxin (LT) (Cohen et al., 1992; Golstein et al., 1991).

The Fas antigen (Fas) is a cell surface protein of relative molecular weight (Mr) 45 kd and carries a single transmembrane domain (Itoh et al., 1991; Oehm et al., 1992; Watanabe-Fukunaga et al., 1992b). Fas is a member of the TNF/nerve growth factor receptor family, which includes two TNF receptors (type I, β or 55 kd; type II, α or 75 kd), the low-affinity nerve growth factor receptor, and CD40, CD27, CD30, and OX40 (Nagata, 1993). Members of this family are homologous as to sequences in the extracellular regions. A high level of Fas mRNA expression has been detected in various tissues, such as the thymus, liver, lung, heart, and ovary of adult mice (Watanabe-Fukunaga et al., 1992b). Some monoclonal antibodies (anti-Fas or anti-AP0-1 antibodies) against human or mouse Fas work as agonists and induce apoptosis of the cells expressing Fas, in vitro and in vivo (Itoh et al., 1991; Ogasawara et al., 1993; Trauth et al., 1989). These results suggested that Fas is a receptor for an unidentified ligand and transduces the apoptotic signal into cells.

The genetic and molecular analyses of the mouse Fas chromosomal gene indicated that Fas is encoded by the gene at the locus of the mouse lymphoproliferation mutation *lpr* (Adachi et al., 1993; Watanabe-Fukunaga et al., 1992a), which is a natural, autosomal, and recessive mutation (Cohen and Eisenberg, 1991). Since mice carrying mutations homozygous at the *lpr* locus have lymphadenopathy and autoimmune disease (Cohen and Eisenberg, 1991), it is likely that Fas is involved in the development of T cells. Accordingly, Fas is expressed in most thymocytes of the wild type, but not in *lpr* mice (Ogasawara et al., 1993). Immature T cells are killed by apoptosis at least in two steps during development in the thymus (Ramsdell and Fowlkes, 1990). Those T cells carrying T cell receptors that do not recognize self MHC antigens as a restriction element are killed or neglected, while the T cells recognizing the self antigens are killed by a process called negative selection. Analysis of thymic T cell development in wild-type and *lpr* mice has suggested that the neglected thymocytes escape from apoptosis in the thymus, then go to the periphery in *lpr* mice (Zhou et al., 1993). In addition to being expressed in thymocytes, Fas is expressed in activated mature T cells (Trauth et al., 1989). Russell et al. (1993) have suggested a role of Fas-mediated apoptosis in the induction of peripheral tolerance, in the antigen-stimulated suicide of mature T cells, or both.

Fas seems to be involved not only in the development and turnover of T cells, but also in cytotoxic T cell (CTL)-mediated apoptosis. Rouvier et al. (1993) showed that the presence of Fas on target cells was required for their lysis by a CTL hybridoma (PC60-d10S [d10S]) and more generally for calcium-independent cytotoxicity by CTLs. These results suggested that CTL cells express the Fas ligand on their surface and that the interaction of Fas ligand with Fas on target cells induces apoptosis in target cells.

To understand the role and mechanism of Fas-mediated apoptosis in various systems, it is essential to identify the Fas ligand. In this study, we isolated a cDNA for the Fas




Figure 1. Flow Cytometry of d10S and COS Cells Transfected with Fas Ligand cDNA

(a and b) Flow cytometry of d10S and d10S-2 cells. The d10S cells were sorted twice by FACS to obtain d10S-2 cells, as described under Experimental Procedures. The d10S cells (1×10^7) (a) or d10S-2 cells (1×10^7) (b) were stained with biotinylated mFas-Fc and PE-conjugated streptavidin before (stippled area) or after (closed area) exposure to PMA and ionomycin for 4 hr. Each sample was washed and analyzed by flow cytometry. The staining profile with PE-conjugated streptavidin alone is also shown (open area).

(c) Flow cytometry of COS cells transfected with pTN24-15. Monkey COS cells (4×10^6) were transfected with pTN24-15. The cells were detached from the plates with EDTA 72 hr later, stained with biotinylated mFas-Fc and PE-labeled streptavidin (closed area), or PE-labeled streptavidin alone (open area), and analyzed by flow cytometry. Control COS cells were transfected with the empty vector, and their staining profile with biotinylated mFas-Fc and PE-conjugated streptavidin is also shown (stippled area).

ligand expressed in d10S cells. The structural analysis of the cDNA indicated that the Fas ligand is a type II membrane protein homologous to members of the TNF family, which includes TNF α , TNF β , LT β , and the ligands for CD40, CD27, and CD30.

Results

Molecular Cloning of the Fas Ligand

To identify the Fas ligand, we prepared a chimeric protein (mFas-Fc) consisting of the extracellular region of the mouse Fas antigen fused to the hinge, CH2, and CH3 domains of human immunoglobulin $\gamma 1$ heavy chain (Aruffo et al., 1990). As a control, a chimeric molecule (hTNFR β -Fc) between the extracellular region of human TNF receptor β (p55) and the Fc region of human immunoglobulin G1 was prepared by a similar method.

The T cell hybridoma, d10S, is cytotoxic against cells expressing Fas, but not against those which do not (Rouvier et al., 1993). The cytotoxic activity of d10S was inhibited by mFas-Fc, but not by hTNFR β -Fc (see below; T. S. and S. N., submitted). When d10S cells were stained with biotinylated mFas-Fc followed by phycoerythrin (PE)-conjugated streptavidin, there was a small shift in a flow cytometric profile (Figure 1a). Stimulation of d10S cells with phorbol myristate acetate (PMA) and ionomycin enhances their cytotoxic activity (Rouvier et al., 1993). Accordingly, this treatment greatly increased the expression of a protein which binds mFas-Fc in d10S cells (Figure 1a). On the other hand, d10S cells were not significantly stained with biotinylated hTNFR β -Fc, indicating that mFas-Fc specifically bound to the Fas ligand (data not shown).

To facilitate cDNA cloning, a population of d10S cells, which expressed large amounts of the Fas ligand, was

selected by FACS using biotinylated mFas-Fc and PE-conjugated streptavidin, as described under Experimental Procedures. To exclude the possible enrichment of Fc γ receptor-positive population, cells were also stained with fluorescein isothiocyanate (FITC)-labeled hTNFR β -Fc, and a cell population (about 0.3%–0.5%) that stained intensely with mFas-Fc but not with hTNFR β -Fc was sorted. Two rounds of sorting of d10S cells (d10S-2) significantly enriched the cell population that highly expressed the Fas ligand, under both unstimulated and stimulated conditions (Figure 1b).

A cDNA library ($\sim 2.2 \times 10^6$ clones) in a mammalian expression vector (pCEV4) (Itoh et al., 1991) was constructed with mRNA from the PMA and ionomycin-stimulated d10S-2 cells. Plasmid DNA from the whole library was introduced into COS cells. The transfected COS cells were allowed to bind mFas-Fc and were then treated with a chemical cross-linker (Harada et al., 1990). The cells cross-linked with mFas-Fc were enriched by panning, as described under Experimental Procedures. This procedure (enrichment by panning and amplification in *Escherichia coli* [E. coli]) was repeated four times, and plasmid DNAs were prepared from individual bacterial clones after the fourth round. Clones (12 out of 48) contained inserts of more than 1.0 kb, and they were individually introduced into COS cells. When these cells were stained with biotinylated mFas-Fc and analyzed by flow cytometry, five of them were positive. A typical result with COS cells transfected with pTN24-15 is shown in Figure 1c. In contrast, COS cells transfected with the empty vector were not stained with biotinylated mFas-Fc. Furthermore, COS cells transfected with pTN24-15 did not significantly bind biotinylated hTNFR β -Fc (data not shown).

Structure of the Fas Ligand

Restriction enzyme mapping of the inserts of the five positive clones indicated that they overlapped. One clone containing a 1.6 kb insert, pTN24-15, was therefore randomly selected and further characterized. The nucleotide sequence of pTN24-15 and its predicted amino acid sequence are presented in Figure 2. The cDNA consists of 1623 nt, and there is only one long open reading frame. The initiation site was tentatively assigned to the ATG codon at nucleotide positions 74–76, although the nucleotide sequence surrounding this initiation site does not conform well with the consensus sequence (CCA/GCCATGG) proposed by Kozak (1991). The open reading frame ends at the termination codon TAA at positions 908–910 and thus codes for a protein of 278 amino acids with a calculated Mr of 31,138 and an isoelectric point of 9.53. The first 77 amino acids of this protein are extremely rich in proline residues. Although it lacks a signal sequence at the N-terminus, its hydropathy analysis indicated that the proline-rich region is followed by 22 hydrophobic amino acids that presumably function as a transmembrane anchor. The lack of a signal sequence and the presence of an internal hydrophobic domain suggested that the Fas ligand is a type II transmembrane protein. The putative extracellular domain in its carboxyl region consists of 179 amino acids and contains four potential N-glycosylation sites (N-X-S/T).

11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65
 66
 67
 68
 69
 70
 71
 72
 73
 74
 75
 76
 77
 78
 79
 80
 81
 82
 83
 84
 85
 86
 87
 88
 89
 90
 91
 92
 93
 94
 95
 96
 97
 98
 99
 100
 101
 102
 103
 104
 105
 106
 107
 108
 109
 110
 111
 112
 113
 114
 115
 116
 117
 118
 119
 120
 121
 122
 123
 124
 125
 126
 127
 128
 129
 130
 131
 132
 133
 134
 135
 136
 137
 138
 139
 140
 141
 142
 143
 144
 145
 146
 147
 148
 149
 150
 151
 152
 153
 154
 155
 156
 157
 158
 159
 160
 161
 162
 163
 164
 165
 166
 167
 168
 169
 170
 171
 172
 173
 174
 175
 176
 177
 178
 179
 180
 181
 182
 183
 184
 185
 186
 187
 188
 189
 190
 191
 192
 193
 194
 195
 196
 197
 198
 199
 200
 201
 202
 203
 204
 205
 206
 207
 208
 209
 210
 211
 212
 213
 214
 215
 216
 217
 218
 219
 220
 221
 222
 223
 224
 225
 226
 227
 228
 229
 230
 231
 232
 233
 234
 235
 236
 237
 238
 239
 240
 241
 242
 243
 244
 245
 246
 247
 248
 249
 250
 251
 252
 253
 254
 255
 256
 257
 258
 259
 260
 261
 262
 263
 264
 265
 266
 267
 268
 269
 270
 271
 272
 273
 274
 275
 276
 277
 278
 279
 280
 281
 282
 283
 284
 285
 286
 287
 288
 289
 290
 291
 292
 293
 294
 295
 296
 297
 298
 299
 300
 301
 302
 303
 304
 305
 306
 307
 308
 309
 310
 311
 312
 313
 314
 315
 316
 317
 318
 319
 320
 321
 322
 323
 324
 325
 326
 327
 328
 329
 330
 331
 332
 333
 334
 335
 336
 337
 338
 339
 340
 341
 342
 343
 344
 345
 346
 347
 348
 349
 350
 351
 352
 353
 354
 355
 356
 357
 358
 359
 360
 361
 362
 363
 364
 365
 366
 367
 368
 369
 370
 371
 372
 373
 374
 375
 376
 377
 378
 379
 380
 381
 382
 383
 384
 385
 386
 387
 388
 389
 390
 391
 392
 393
 394
 395
 396
 397
 398
 399
 400
 401
 402
 403
 404
 405
 406
 407
 408
 409
 410
 411
 412
 413
 414
 415
 416
 417
 418
 419
 420
 421
 422
 423
 424
 425
 426
 427
 428
 429
 430
 431
 432
 433
 434
 435
 436
 437
 438
 439
 440
 441
 442
 443
 444
 445
 446
 447
 448
 449
 450
 451
 452
 453
 454
 455
 456
 457
 458
 459
 460
 461
 462
 463
 464
 465
 466
 467
 468
 469
 470
 471
 472
 473
 474
 475
 476
 477
 478
 479
 480
 481
 482
 483
 484
 485
 486
 487
 488
 489
 490
 491
 492
 493
 494
 495
 496
 497
 498
 499
 500
 501
 502
 503
 504
 505
 506
 507
 508
 509
 510
 511
 512
 513
 514
 515
 516
 517
 518
 519
 520
 521
 522
 523
 524
 525
 526
 527
 528
 529
 530
 531
 532
 533
 534
 535
 536
 537
 538
 539
 540
 541
 542
 543
 544
 545
 546
 547
 548
 549
 550
 551
 552
 553
 554
 555
 556
 557
 558
 559
 560
 561
 562
 563
 564
 565
 566
 567
 568
 569
 570
 571
 572
 573
 574
 575
 576
 577
 578
 579
 580
 581
 582
 583
 584
 585
 586
 587
 588
 589
 590
 591
 592
 593
 594
 595
 596
 597
 598
 599
 600
 601
 602
 603
 604
 605
 606
 607
 608
 609
 610
 611
 612
 613
 614
 615
 616
 617
 618
 619
 620
 621
 622
 623
 624
 625
 626
 627
 628
 629
 630
 631
 632
 633
 634
 635
 636
 637
 638
 639
 640
 641
 642
 643
 644
 645
 646
 647
 648
 649
 650
 651
 652
 653
 654
 655
 656
 657
 658
 659
 660
 661
 662
 663
 664
 665
 666
 667
 668
 669
 670
 671
 672
 673
 674
 675
 676
 677
 678
 679
 680
 681
 682
 683
 684
 685
 686
 687
 688
 689
 690
 691
 692
 693
 694
 695
 696
 697
 698
 699
 700
 701
 702
 703
 704
 705
 706
 707
 708
 709
 710
 711
 712
 713
 714
 715
 716
 717
 718
 719
 720
 721
 722
 723
 724
 725
 726
 727
 728
 729
 730
 731
 732
 733
 734
 735
 736
 737
 738
 739
 740
 741
 742
 743
 744
 745
 746
 747
 748
 749
 750
 751
 752
 753
 754
 755
 756
 757
 758
 759
 760
 761
 762
 763
 764
 765
 766
 767
 768
 769
 770
 771
 772
 773
 774
 775
 776
 777
 778
 779
 780
 781
 782
 783
 784
 785
 786
 787
 788
 789
 790
 791
 792
 793
 794
 795
 796
 797
 798
 799
 800
 801
 802
 803
 804
 805
 806
 807
 808
 809
 810
 811
 812
 813
 814
 815
 816
 817
 818
 819
 820
 821
 822
 823
 824
 825
 826
 827
 828
 829
 830
 831
 832
 833
 834
 835
 836
 837
 838
 839
 840
 841
 842
 843
 844
 845
 846
 847
 848
 849
 850
 851
 852
 853
 854
 855
 856
 857
 858
 859
 860
 861
 862
 863
 864
 865
 866
 867
 868
 869
 870
 871
 872
 873
 874
 875
 876
 877
 878
 879
 880
 881
 882
 883
 884
 885
 886
 887
 888
 889
 890
 891
 892
 893
 894
 895
 896
 897
 898
 899
 900
 901
 902
 903
 904
 905
 906
 907
 908
 909
 910
 911
 912
 913
 914
 915
 916
 917
 918
 919
 920
 921
 922
 923
 924
 925
 926
 927
 928
 929
 930
 931
 932
 933
 934
 935
 936
 937
 938
 939
 940
 941
 942
 943
 944
 945
 946
 947
 948
 949
 950
 951
 952
 953
 954
 955
 956
 957
 958
 959
 960
 961
 962
 963
 964
 965
 966
 967
 968
 969
 970
 971
 972
 973
 974
 975
 976
 977
 978
 979
 980
 981
 982
 983
 984
 985
 986
 987
 988
 989
 990
 991
 992
 993
 994
 995
 996
 997
 998
 999
 1000

Figure 2. Nucleotide Sequence and Predicted Amino Acid Sequence of Rat Fas Ligand cDNA. The numbers above and below each line refer to the nucleotide position and amino acid positions, respectively. The putative transmembrane domain is underlined and four potential N-linked glycosylation sites (N-X-S/T) are indicated by asterisks.

The d10S cell line used to prepare the cDNA library is a hybrid between rat and mouse cell lines (Conzelmann et al., 1982). To determine the origin of the cloned Fas ligand cDNA, two sets of oligonucleotides containing sequences for the 3' noncoding region of pTN24-15 (nucleotides from positions 1006-1025 and 1305-1324 for the forward primer and 1327-1346 and 1543-1562 for the reverse primer) were synthesized. When genomic DNAs from the rat and mouse spleen were analyzed by polymerase chain reaction, only rat chromosomal DNA gave bands of the predicted sizes of 341 bp and 258 bp (data not shown). Furthermore, Northern hybridization with pTN24-15 as probe under high stringency gave a band of 2 kb in rat but not mouse RNA (see below, T. T., S. S., and S. N., unpublished data). These results indicated that the cloned cDNA sequence was derived from the rat genome present in the d10S hybrid cell line.

Biochemical and Biological Properties of the Fas Ligand

The Fas ligands expressed in d10S cells and in the COS cells transfected with pTN24-15 were biochemically characterized by immunoprecipitation using mFas-Fc. The mFas-Fc specifically precipitated protein(s) of Mr 38-42 kd from biotinylated cell surface proteins of the d10S-12 cell line (sorted 12 times by FACS) (Figure 3; T. S. and S. N., submitted). Similarly, immunoprecipitation of the lysates from COS cells transfected with pTN24-15, but not with the empty vector, gave several bands of Mr 36-43 kd, which differed slightly from the bands detected in d10S-12

cells. The Mr of the Fas ligand expressed in d10S-12 cells or COS cells is larger than that calculated from its amino acid sequence (Mr, 31,138). The difference is probably due to glycosylation of some of the four N-glycosylation sites. This may also explain the slightly different Mr between the natural Fas ligand in d10S cells and the recombinant Fas ligand expressed in COS cells.

To confirm that pTN24-15 codes for the Fas ligand, the cytotoxic activity of the COS cells expressing the recombinant Fas ligand was examined using WR19L transformants (W4) that express mouse Fas (Ogasawara et al., 1993) as target cells. As shown in Figure 4b, COS cells transfected with pTN24-15 induced cytotoxicity of W4 in a dose-dependent manner, while control COS cells transfected with the empty vector exhibited no cytotoxic activity. The transfected COS cells were at least 10 times more efficient as effectors against W4 cells, based upon the effector/target cells ratio, than were the original d10S cells (Figures 4a and 4b). The Fas ligand expressed in d10S or COS cells could not kill parental WR19L cells (Figure 4). Since WR19L cells are susceptible to killing by TNF (Itoh et al., 1993), this suggests that the Fas ligand cannot induce the death signal through the TNF receptor. To confirm the specificity of the Fas ligand, mFas-Fc or hTNFR β -Fc was added to the assay mixture. As shown in Figure 4d, the cytotoxic activities of COS cells expressing the recombinant Fas ligand, as well as that of the d10S cells, were almost completely inhibited by 10 μ g/ml of mFas-Fc, but not by hTNFR β -Fc. The supernatant of COS cells transfected with the Fas ligand cDNA also had significant




Figure 3. Immunoprecipitation of the Fas Ligand with the mFas-Fc. The d10S cells were sorted 12 times by FACS to obtain the d10S-12 cell line. COS cells were transfected with pCEV4 vector DNA or pTN24-15 carrying Fas ligand cDNA. After 72 hr, transfected COS cells were detached with EDTA. The cell surface proteins of the d10S-12 cell line and the transfected COS cells were biotinylated, lysed in buffer containing NP-40, precleared with hTNFR8-Fc, and immunoprecipitated with mFas-Fc, as described under Experimental Procedures. The immunoprecipitates were heated at 95°C for 2 min in Laemmli's sample buffer containing 5% 2-mercaptoethanol and resolved by electrophoresis through a 10%–20% gradient polyacrylamide gel in the presence of 0.1% SDS. The proteins were blotted onto a PVDF membrane (Millipore) and detected using the ECL system (Amersham) after staining with streptavidin-conjugated horseradish peroxidase. As size markers, biotin-labeled molecular weight standards (Pharmacia) were electrophoresed in parallel; sizes of standard proteins are shown in kilodaltons.

cytotoxic activity upon W4, but not upon WR19L parental cells (Figure 4c). These results suggest that the recombinant Fas ligand expressed in COS cells can be released into the extracellular fluid, perhaps by cleavage from the cell surface.

We and others have reported that the Fas activated by agonistic anti-Fas antibodies mediates apoptosis [Itoh et al., 1991; Ogasawara et al., 1993; Trauth et al., 1989]. The recombinant Fas ligand expressed on COS cells also induced apoptosis. In Figure 5, the chromosomal DNA was prepared from W4 cells after various periods of incubation with COS cells that were transfected either with pTN24-15 or with the empty vector pCEV4. COS cells transfected with pTN24-15 induced fragmentation of the chromosomal DNA of W4 cells in a step ladder fashion that is characteristic of apoptosis [Compton, 1992]. The DNA ladder was





Figure 4. Cytotoxic Activity of d10S Cells and the Recombinant Fas Ligand Expressed in COS Cells

(a) Cytotoxic activity of d10S cells. The cytotoxic activity was assayed in duplicate using ^{51}Cr -labeled W4 (closed circle) or WR19L cells (open circle), as described under Experimental Procedures. (b) Cytotoxic activity of the recombinant Fas ligand expressed in COS cells. COS cells were transfected with the pCEV4 vector (closed circle) or pTN24-15 DNA (closed triangle, open circle). At 72 hr after transfection, cytotoxicity of the transfected COS cells using W4 (closed circle, closed triangle) or WR19L cells (open circle) as target cells is shown. (c) Cytotoxic activity of the soluble recombinant Fas ligand produced by COS cells. COS cells were transfected with pCEV4 (closed circle) or pTN24-15 (closed triangle, open circle). At 18 hr after transfection, the medium was changed to serum-free DMEM, and cells were incubated for 48 hr. The culture supernatant of the transfected cells was filtered through 0.22 μm filter, concentrated 8-fold, and then added to the target cells. W4 cells (closed circle, closed triangle) or WR19L cells (open circle) were used as target cells. One-fold concentration corresponds to the neat supernatant of COS cells. (d) Inhibition of the cytotoxic activity of the Fas ligand with mFas-Fc. The cytotoxic activity of the activated d10S cells at an effector/target (E/T) ratio of 10 (lanes 1–3) or COS cells transfected with pTN24-15 at an E/T ratio of 0.2 (lanes 4–6) was determined using ^{51}Cr -labeled W4 cells as target cells. No chimeric proteins (open bar), 10 $\mu\text{g}/\text{ml}$ of hTNFR8-Fc (hatched bar), or mFas-Fc (closed bar) was added to the reaction mixture.

observed within 1 hr of incubation, and most of it was degraded into nucleosome-sized fragments after 2 hr. Such fragmentation was not observed in W4 cells incubated with COS cells transfected with pCEV4 (Figure 5). Moreover, the activity of the recombinant Fas ligand was seen only in W4 cells, but not in WR19L cells (data not shown).



Figure 5. DNA Fragmentation Induced by the Recombinant Fas Ligand. COS cells were transfected with pCEV4 vector (lanes 1–4) or the Fas ligand expression plasmid pTN24-15 (lanes 5–9). At 48 hr after transfection, W4 cells were added to the transfected COS cells and incubated at 37°C. Before incubation (lanes 1 and 9) and after incubation for 1 hr (lanes 2 and 7), 2 hr (lanes 3 and 8), or 3 hr (lanes 4 and 6), total DNA was prepared from the nonadherent cells and resolved on a 1.0% agarose gel in the presence of 0.5 $\mu\text{g}/\text{ml}$ of ethidium bromide. For size markers (lane 5), BamHI-digested and HinfI-digested pBR322 was electrophoresed in parallel; sizes of the DNA fragments are given in base pairs.



The original and sorted d10S cell lines, rat splenocytes and thymocytes, and various rat tissues were examined by Northern blot hybridization using the Fas ligand cDNA as a probe. As shown in Figure 6A, poly(A) RNA from the original d10S cells showed a weakly hybridizing band of 2 kb. The intensity of this band increased considerably after the d10S cells were activated with PMA and ionomycin. The d10S-2 cell line, which was sorted twice by FACS, expressed about four times more Fas ligand mRNA than did the original d10S after stimulation with PMA and ionomycin, which is consistent with the results of the flow cytometry (see Figure 1). Recently, we obtained a d10S-16 cell line (sorted 16 times) that exhibited about 100 times higher staining intensity with mFas-Fc and 100 times higher cytotoxic activity than did the original d10S cells (T. S. and S. N., submitted). Accordingly, d10S-16 cells had about 100 times more Fas ligand mRNA than did the original d10S cells (Figure 6A). Thus, the increased expression of Fas ligand mRNA correlated well with the increased staining intensity with mFas-Fc and the cytotoxic activity.

Discussion

or after (plus sign) stimulation with PMA and ionomycin. One microgram of mRNA per lane was electrophoresed on a 1.5% agarose gel and analyzed by Northern hybridization using the ³²P-labeled Fas ligand cDNA (Fas-L, upper panel) or human EF-1α cDNA (lower panel). The positions of 18S and 28S rRNAs are indicated on the left.

(B) Expression of the Fas Ligand in Rat Splenocytes and Thymocytes. Freshly isolated rat splenocytes or thymocytes were cultured at 37°C for 8 hr in the medium alone or in that containing 10 ng/ml PMA and 500 ng/ml ionomycin or 5 µg/ml concanavalin A and 10 ng/ml human interferon 2. Poly(A) RNA, 2.0 µg per lane, was subjected to Northern hybridization with the rat Fas ligand cDNA (Fas-L, upper panel) or human EF-1α cDNA as a probe (lower panel).

(C) Expression of the Fas Ligand mRNA in Rat Tissues. Poly(A) RNA was prepared from the indicated rat tissues, and 2.0 µg RNA per lane was analyzed by Northern hybridization using rat Fas ligand cDNA (Fas-L, upper panel) and human EF-1α cDNA (lower panel) as probe.




Figure 7. Alignment of the Amino Acid Sequence of the TNF Family Members

The C-terminal amino acid sequence of the rat Fas ligand was aligned with the corresponding amino acid sequences of the other members of the TNF family, beginning at the N-terminus of mature soluble human TNF α . Numbers in brackets indicate the positions from the N-terminus of the primary translation products. Several gaps have been introduced to optimize the alignment. Amino acids identical in more than five members are indicated by asterisks. The amino acids of favored substitutions in more than four members are shown in bold letters. Favored amino acid substitutions are defined as pairs of residues belonging to one of the following groups: S, T, P, A, and G; N, D, E, and Q; H, R, and K; M, I, L, and V; and F, Y, and W. The extent of the β sheet regions a-h of TNF α (Eck and Sprang, 1989) is shown at the top of each line. The cysteine residues found in the corresponding positions of the Fas ligand and TNF α are underlined.

of apoptosis also seem to be regulated by cytokines and their receptors (Nagata, 1993). We previously suggested that Fas is a receptor for an unknown death factor and mediates apoptosis. In this report, the cDNA for the death factor, or the Fas ligand, was isolated from a CTL cell line.

The Fas belongs to the TNF/nerve growth factor receptor family, which includes two TNF receptors, the low affinity nerve growth factor receptor, and CD40, CD30, and CD27 (Itoh et al., 1991; Nagata, 1993). TNF α exists either as a type II transmembrane protein or a soluble cytokine (Kriegler et al., 1988). The ligands for CD40, CD27, and CD30 have been identified as type II transmembrane proteins related to TNF (Armitage et al., 1992; Smith et al., 1993; Goodwin et al., 1993). Recently, it was demonstrated that LTP, which can associate with TNF β (LT α) on the cell surface, is also a member of the TNF family (Browning et al., 1993). The amino acid sequence of the Fas ligand is clearly homologous to those members of the TNF family (Figure 7). Homologous regions are restricted to the carboxyl region; that is, the extracellular domain that interacts with the receptor. A pairwise comparison of the amino acid sequences in this region indicated that the Fas ligand is more similar to TNF α , TNF β , and LT β than to ligands for CD40, CD27, and CD30. The high homologies of the Fas ligand with TNF α and TNF β (27.0% and 28.7% identity, respectively) are comparable to that between TNF α and TNF β (29.6% identity) (Pennica et al., 1984). Despite this similarity, the cloned rat Fas ligand

could neither bind to the 55 kd human TNF receptor nor activate the mouse TNF receptor in WR19L cells.

TNF α and TNF β function as trimers (Smith and Baglioni, 1987). X-ray diffraction analyses of TNF α and TNF β have indicated that each monomer of these cytokines forms an elongated, antiparallel β -pleated sheet sandwich with a jelly roll topology (Eck and Sprang, 1989; Eck et al., 1992; Jones et al., 1989; Banner et al., 1993). As shown in Figure 7, the conserved amino acids among the Fas ligand, TNF α , and TNF β were found mainly within the β strands, suggesting the Fas ligand has a similar structure to TNF α and TNF β . Furthermore, in TNF α , a disulfide bridge links the loop connecting β strands c and d with that connecting β strands e and f (Eck and Sprang, 1989). The two cysteine residues in the extracellular domain of the Fas ligand were found at the corresponding positions (Figure 7), suggesting that these cysteine residues are connected by a disulfide bond similar to that found in TNF α . These structural similarities of the Fas ligand with TNF α suggest that the Fas ligand also exists as a trimer. The original anti-Fas or anti-APO-1 antibody works as an agonist that induces apoptosis by binding to Fas (Itoh et al., 1991; Trauth et al., 1989). The anti-Fas antibody is an immunoglobulin M (Yonehara et al., 1989), which is an immunoglobulin pentamer, whereas the anti-APO-1 antibody is an immunoglobulin G α (Trauth et al., 1989), which tends to aggregate. The F(ab) $_2$ fragment of the anti-APO-1 antibody and its isotypes have little cytotoxic activity (Dhein et al., 1992). However, these divalent anti-APO-1 antibodies induced apoptosis when they were cross-linked by second antibodies. These results indicate that dimerization of Fas is insufficient to transduce the apoptotic signal, and they are consistent with a trimeric structure for the Fas ligand.

Although Rouvier et al. (1993) could not detect Fas ligand activity in the culture medium of the d10S cell line, we detected Fas-dependent cytotoxic activity in the supernatant of COS cells transfected with a Fas ligand expression plasmid. Moreover, the medium conditioned with the d10S-16 cells, which expresses about 100 times more Fas ligand mRNA than the original d10S cell line (Figure 6A), was significantly cytotoxic against cells expressing Fas (T. S. and S. N., unpublished data). These results suggest that overexpression of the Fas ligand on the cell surface causes shedding of the protein, as found in TNF (Kriegler et al., 1988; Perez et al., 1990), and the soluble form of the Fas ligand actively triggers apoptosis by binding to Fas. A large amount of soluble TNF is present in the serum of individuals suffering from septic shock caused by bacterial endotoxins or in the serum of cancer patients (Beutler and Cerami, 1986; Old, 1987). Similarly, production of the soluble Fas ligand may accompany some human diseases. Since administration of agonistic anti-Fas antibody into mice causes a phenotype similar to that of fulminant hepatitis (Ogasawara et al., 1993), it will be of interest to examine the involvement of the soluble Fas ligand in human fulminant hepatitis.

Various effector molecules are thought to be involved in CTL-mediated cytotoxicity (Apasov et al., 1993; Golstein et al., 1991; Podack et al., 1991). One well characterized pathway is the perforin-mediated cytotoxicity of target cells,

which is dependent on Ca^{2+} . It is also suggested that the soluble or membrane-bound form of TNF or LT is involved in various target cell systems. Recently, Rouvier et al. (1993) demonstrated that the cytotoxic activity of CTL hybridoma d10S depends on the expression of Fas on target cells. Here, we confirmed by generating cytotoxicity with the recombinant Fas ligand that this cytotoxicity is mediated by interaction between the Fas ligand on d10S cells and Fas on target cells. Furthermore, the cytotoxic activity of transfectants of COS (a fibroblast-like kidney-derived cell line) strongly argues that the cytotoxicity mediated by the Fas-Fas ligand system requires no other molecules preferentially expressed in activated lymphocytes (such as perforin or granzymes). Similar Fas-dependent cytotoxicity, which is alloantigen-specific and Ca^{2+} -independent, has been observed in peritoneal exudate lymphocytes (Rouvier et al., 1993). It is likely that the interaction of the relevant MHC on target cells with the T cell receptor on effector cells induces the Fas ligand on the latter, which then bind to Fas on target cells to induce apoptosis (Rouvier et al., 1993). In addition, we detected Fas ligand mRNA in rat splenocytes after stimulation with PMA and ionomycin, or concanavalin A and interleukin 2. These results suggest that the Fas system is a novel and common mechanism of T cell-mediated cytotoxicity. In this regard, it is noteworthy that some $CD4^+$ Th1 cell lines exhibit a cytolytic activity independent of perforin and TNF (Ju, 1991; Ozdemirli et al., 1992). It would be of interest to examine whether or not the Fas system is also involved in these cytolytic activities.

Mice carrying the *lpr* or *gld* mutation develop lymphadenopathy and autoimmune disease (Cohen and Eisenberg, 1991). It has been suggested that the *lpr* and *gld* are mutations of genes encoding an interacting pair of molecules involved in T cell development (Allen et al., 1990). We have previously demonstrated that *lpr* mice have defects in the Fas gene and thus proposed that the Fas ligand is encoded by the gene at the *gld* locus (Watanabe-Fukunaga et al., 1992a). Recent findings that peritoneal exudate lymphocytes from *gld* mice do not show the Fas-dependent cytotoxicity (P. G., unpublished data) support our notion. To confirm this theory, it would be necessary to correlate the chromosomal gene locus of mouse Fas ligand with the *gld* locus and to examine the abnormality of the chromosomal gene for the Fas ligand in *gld* mice. The *lpr* or *gld* mice are thought to have defects in the thymic development of T cells (Cohen and Eisenberg, 1991). We previously suggested that the expression of the Fas in thymocytes and the Fas ligand in thymic stromal cells control the development of T cells (Watanabe-Fukunaga et al., 1992a). Although most thymocytes express Fas (Ogasawara et al., 1993), we could not detect the Fas ligand transcript in the thymus except for weak expression in activated thymocytes. It is possible that the Fas ligand is expressed only transiently in a limited number of stromal cells in the thymus. The availability of Fas ligand cDNA and mFas-Fc fusion protein would facilitate the identification of such cells by *in situ* hybridization and immunohistochemistry.

The Fas is expressed not only in the cells of the immune

system, but also in the liver, lung, ovary, and heart (Watanabe-Fukunaga et al., 1992b). The physiological functions of Fas in these tissues are not clear, and no developmental abnormalities of these tissues have been noticed in *lpr* or *gld* mice. There was no major expression of Fas ligand mRNA in the aforementioned tissues, although, for instance, the lung expresses Fas and some detectable level of Fas ligand mRNA. On the other hand, the Fas ligand was expressed in the testis, where no apparent expression of Fas was previously detected (Watanabe-Fukunaga et al., 1992b). The testicular seminiferous epithelium is highly proliferative, and many germ cells are programmed to die by apoptosis by interaction with Sertoli cells (Allan et al., 1992; Miething, 1992). Whether the Sertoli cells express the Fas ligand to kill the germ cells remains to be examined. In any event, these results are consistent with the involvement of the Fas system in various aspects of mammalian development.

Experimental Procedures

Flow Cytometry and Selection of the d10S-2 Cell Line

The expression plasmids for mFas-Fc and hTNFR β -Fc were constructed with mouse Fas gene (Watanabe-Fukunaga et al., 1992b) or human TNFR expression plasmid p55TNFR-HO1 (Loetscher et al., 1991) and human immunoglobulin gene in plasmid pMH4 (Nishimura et al., 1987). These chimeric proteins were transiently produced in COS cells or in stable transformants of BTS-1 cells (Sedivy et al., 1988) and were homogeneously purified. The details of the production procedure will be described elsewhere (T. S. and S. N., submitted). The mFas-Fc and hTNFR β -Fc were biotinylated using sulfoacetyl-*N*-hydroxysuccinimide (NHS-LC-biotin, Pierce) according to the protocol of the manufacturer. To prepare FITC-conjugated human TNFR β -Fc, 1 mg of protein was mixed with 20 μ g of FITC in 1 ml of 50 mM sodium carbonate buffer (pH 9.5). After incubation at room temperature for 4 hr, free FITC was removed by Sephadex G-25M column chromatography. For flow cytometry, d10S or transfected COS cells were washed with staining solution (phosphate-buffered saline [PBS] containing 2% fetal calf serum [FCS] and 0.02% NaN₃). Cells (1×10^6) were first incubated on ice for 10 min in 50 μ l of staining solution containing 5 μ g/ml rat anti-mouse Fc γ R receptor antibody (Pharmingen). Fifty microliters of biotinylated mFas-Fc (10 μ g/ml) was added to the reaction mixture and incubated on ice for 30 min. After washing with staining solution, the cells were stained on ice for 30 min with PE-conjugated streptavidin (25-fold dilution, Becton-Dickinson) in 100 μ l of staining solution. Cells were washed with staining solution and analyzed by flow cytometry with a FACScan (Becton-Dickinson).

A subline of the d10S cell line that stained intensely with mFas-Fc was selected by repetitive FACS sorting. In brief, 1×10^6 to 3×10^6 d10S cells were stained with FITC-conjugated hTNFR β -Fc and biotinylated mFas-Fc followed by PE-conjugated streptavidin, as described above, and sorted using a FACStar (Becton-Dickinson). The cells providing the highest levels of PE-fluorescence signal (top 0.3%–0.5%), but not significantly stained by FITC-hTNFR β -Fc, were collected and expanded in Dulbecco's modified Eagle's medium (DMEM) containing 10% FCS and 50 mM 2-mercaptoethanol.

Cytotoxicity Assays

Cytotoxicity was assayed essentially as described previously (Rouvier et al., 1993). WR19L or W4 cells (1×10^6) were incubated for 2 hr at 37°C with 20 μ Ci of [³H] sodium chromate (Amersham) in 100 μ l of RPMI1640 containing 10% FCS. After washing with medium, these cells were used as the target. Transfection of COS cells with the pTN24-15 or the control vector was performed by DEAE-dextran method, as described previously (Fukunaga et al., 1990). The [³H]-labeled target cells (1×10^6) were mixed with original d10S cells or transfected COS cells at various ratios in round-bottomed microtiter plates in a total volume of 200 μ l. The plates were centrifuged at 700 rpm for 2 min and incubated for 4 hr at 37°C. The plates were then

centrifuged at 1200 rpm for 5 min, and 100 μ l aliquots of the supernatants were assayed for radioactivity using a γ -counter. The spontaneous release of ^{51}Cr was determined by incubating the target cells with the medium alone, whereas the maximum release was determined by adding Triton X-100 to a final concentration of 0.1%. The percentage of specific ^{51}Cr release was calculated as follows:

$$\% \text{ specific lysis} = \frac{\text{experimental } ^{51}\text{Cr release} - \text{spontaneous } ^{51}\text{Cr release}}{\text{maximum } ^{51}\text{Cr release} - \text{spontaneous } ^{51}\text{Cr release}}$$

The spontaneous release of ^{51}Cr was routinely 8%–10% of the maximum release.

Construction of the cDNA Library

The d10S-2 cells were grown up to 2×10^6 cells/ml in DMEM containing 10% FCS and were then stimulated with 20 ng/ml PMA and 1 μ g/ml ionomycin at 37°C for 3 hr. Total cellular RNA was isolated using guanidine isothiocyanate-phenol acid (Chomczynski and Sacchi, 1987), and poly(A) RNA was selected by two cycles of oligo(dT)-cellulose column chromatography. Double-stranded cDNA primed with a random hexamer or oligo(dT) was synthesized as described (Itoh et al., 1991). A BstOI adaptor was added to the cDNA, which was size-fractionated by electrophoresis on a 1% agarose gel. Those cDNAs larger than 1.5 kb were recovered and ligated to BstOI-digested pCEV4. E. coli DH10B cells (GIBCO BRL) were transformed with the ligated DNA by electroporation (Dower et al., 1988). About 1.0×10^6 independent clones from an oligo(dT)-primed cDNA library were mixed with 1.3×10^6 clones from a random hexamer-primed cDNA library and were used to transfect COS7 cells.

Enrichment of cDNA Clones for the Fas Ligand by Panning

COS7 cells were transfected with plasmid DNA by electroporation (Potter et al., 1994). In brief, 5×10^6 COS cells were washed with K-PBS* (30.8 mM NaCl, 120.7 mM KCl, 8.1 mM Na_2HPO_4 , and 1.46 mM KH_2PO_4), and suspended in 0.4 ml of K-PBS* supplemented with 5 mM MgCl_2 (K-PBS*). Plasmid DNA (40 μ g) dissolved in 0.4 ml of K-PBS* was added to the cell suspension and incubated on ice for 10 min. Cells were then exposed to a 230 V pulse with a capacitance of 960 μ F and returned to the ice. After 10–15 min, the cell suspensions were diluted with 5 ml of cold serum-free DMEM and incubated for 30 min at room temperature. The cells were then plated on 2 10-cm dishes and cultured for 80 hr at 37°C in DMEM containing 10% FCS. A total of 1.2×10^6 COS cells were transfected as above and cultured in 96 10-cm plates. The cells were detached from the plates by incubating at 37°C for 30 min in 5 ml per plate of PBS-EDTA- NaN_3 (PBS containing 0.5 mM EDTA and 0.02% NaN_3). The detached cells were resuspended to a concentration of 5×10^6 to 7×10^6 cells per milliliter in PBS-EDTA- NaN_3 containing 3 mg/ml of bovine serum albumin and 2.5 μ g/ml anti-mouse Fc γ 1 receptor antibody. After incubation on ice for 10 min, mFas-Fc was added to a final concentration of 4 μ g/ml and incubated on ice for 60 min. The cells were washed with ice-cold PBS and suspended to 5×10^6 to 7×10^6 cells per milliliter in PBS containing 50 mM HEPES buffer (pH 8.3) and 0.2 mM bis(sulfosuccinimidyl) suberate (BS 2 , Pierce). After incubation on ice for 30 min, 1 M Tris-HCl (pH 8.0) was added to a final concentration of 50 mM and incubated on ice for 10 min. After washing with PBS, cells were suspended in 30 ml of PBS-EDTA- NaN_3 containing 3 mg/ml bovine serum albumin and passed through Nylon mesh (pore size, 100 μ m) to remove aggregates. The cells were then distributed into 30 10-cm panning plates that were coated with goat anti-human immunoglobulin G Fc (Cappel). After incubation for 2 hr at room temperature, nonadherent cells were removed by gentle washing with PBS-EDTA- NaN_3 containing 3 mg/ml bovine serum albumin, and the extrachromosomal DNA was extracted from the adherent cells as described (Itoh et al., 1991). Electroporation of E. coli with the DNA obtained from the first round of panning yielded 4.1×10^6 colonies, from which plasmid DNA was prepared, and was used to transfect 9.8×10^6 COS cells (50 10-cm plates). The second panning using 30 plates and DNA preparation from the adherent cells proceeded as described above. Transformation of E. coli with the recovered plasmid DNA yielded 8×10^6 clones, and the third panning using 30 plates was performed by transfecting 4×10^6 COS cells (in 10 plates). With the plasmid DNA recovered from the adherent cells, E. coli was transformed to produce 3.8×10^6

clones, which were subjected to a fourth panning, using 1×10^6 COS cells (in 25 plates) and 10 panning plates.

DNA Sequence Analysis and Northern Hybridization

The DNA sequence was determined using a DNA sequencer (model 370A; Applied Biosystems) and a Taq DyeDeoxy cycle sequencing kit from Applied Biosystems. For Northern hybridization, poly(A) RNA was prepared from d10S and its derivatives and various rat tissues and cells using a mRNA isolation kit from Pharmacia. RNA was denatured at 65°C for 5 min in 50% formamide, electrophoresed through a 1.5% agarose gel containing 8.8% formaldehyde, and transferred to nitrocellulose or nylon membranes (Schleicher & Schuell). The probe was a 0.9 kb DNA fragment containing the sequence from 43 to 967 of pTN24-15, prepared by PCR, and labeled with ^{32}P using a random primer labeling kit (Boehringer, Mannheim). A ^{32}P -labeled 1.8 kb BamHI fragment of human EF-1 α cDNA (Uetsuki et al., 1989) was used as a control probe DNA. Hybridization proceeded as described (Sambrook et al., 1989) under high stringency.

Fragmentation of Chromosomal DNA

COS cells (8×10^7) in 24 well plates were transfected with 1.0 μ g of pTN24-15 using DEAE-dextran as described (Fukunaga et al., 1990). WR19L cells or W4 cells (Opasawara et al., 1993) (2×10^7) were added to the wells 48 hr after transfection and incubated in RPMI1640 medium containing 10% FCS at 37°C. Nonadherent cells were collected, and chromosomal DNA was prepared as described (Laird et al., 1991).

Surface Labeling and Immunoprecipitation

The cell surface proteins of d10S-12 cells or COS cells transfected with pTN24-15 were biotinylated using D-biotinyl-L-aminocaproic acid N-hydroxysuccinimide ester (biotin-CNHS-ester, Boehringer-Mannheim) as described (Meier et al., 1992). The cells (7.5×10^6 cells) were lysed by incubating on ice for 30 min in 1 ml of lysis buffer (1% NP-40, 50 mM Tris-HCl (pH 8.0), 150 mM NaCl, 1 mM D-aminophenylmethanesulfonyl fluoride hydrochloride [APMSF], 1 μ g/ml pepstatin, and 1 μ g/ml leupeptin). After centrifugation at 14,000 rpm for 15 min, the supernatant was precleared by incubation on ice for 60 min with 10 μ g/ml of hTNFR β -Fc and then at 4°C for 60 min with a 5% volume of protein A-Sepharose. After removing the protein A-Sepharose, 10 μ g of mFas-Fc was added to the supernatant and incubated on ice for 60 min. Protein A-Sepharose (3.01 vol) was added to the mixture and incubated at 4°C overnight. After centrifugation, the precipitates were washed with the lysis buffer and resuspended in 20 μ l of Laemmli's sample buffer (62.5 mM Tris-HCl (pH 6.8), 2% SDS, 10% glycerol, and 0.002% bromophenol blue).

Acknowledgments

We thank Dr. T. Watanabe for plasmid pMH4, Dr. W. Lesslauer for plasmid p55TNF-HG1, and Dr. J. M. Sedivy for BST-1 cells. We are grateful to Dr. N. Roh for advice concerning the panning procedure and Ms. K. Mimura for secretarial assistance. This work was supported in part by grants-in-aid from the Ministry of Education, Science, and Culture of Japan.

Received October 4, 1993; revised October 22, 1993.

References

- Adachi, M., Watanabe-Fukunaga, R., and Negata, S. (1993). Aberrant transcription caused by the insertion of an early transposable element in an intron of the Fas antigen gene of *lpr* mice. *Proc. Natl. Acad. Sci. USA* 90, 1758–1760.
- Altan, D. J., Harmon, B. V., and Roberts, S. A. (1992). Spermatogenic apoptosis has three morphologically recognizable phases and shows no circadian rhythm during normal spermatogenesis in the rat. *Cell Prolif.* 25, 241–250.
- Allen, R. D., Marshall, J. D., Roths, J. B., and Sidman, C. L. (1990). Differences defined by bone marrow transplantation suggest that *lpr* and *gld* are mutations of genes encoding an interacting pair of molecules. *J. Exp. Med.* 172, 1367–1375.

- Apasov, S., Redegeld, F., and Sitarovsky, M. (1993). Cell-mediated cytotoxicity: contact and secreted factors. *Curr. Opin. Immunol.* 5, 404-410.
- Armitage, R. J., Fanslow, W. C., Strockbine, L., Sato, T. A., Clifford, K. N., Macduff, B. M., Anderson, D. M., Gimpel, S. D., Davis-Smith, T., Maliszewski, C. R., Clark, E. A., Smith, C. A., Grabstein, K. H., Cosman, D., and Spriggs, M. K. (1992). Molecular and biological characterization of a murine ligand for CD40. *Nature* 357, 80-82.
- Aruffo, A., Stamenkovic, I., Melnick, M., Underhill, C. B., and Seed, B. (1990). CD44 is the principal cell surface receptor for hyaluronate. *Cell* 61, 1303-1313.
- Banner, D. W., D'Arcy, A., Jones, W., Gentz, R., Schoenfeld, H.-J., Broger, C., Loetscher, H., and Lesslauer, W. (1993). Crystal structure of the soluble human 55 kD TNF receptor-human TNF complex: implications for TNF receptor activation. *Cell* 73, 431-445.
- Beutler, B., and Cerami, A. (1986). Cachectin and tumor necrosis factor as two sides of the same biological coin. *Nature* 320, 584-588.
- Browning, J. L., Ngam-ek, A., Lawton, P., DeMarinis, J., Tizard, R., Chow, E. P., Hession, C., O'Brien-Greco, B., Foley, S. F., and Ware, C. F. (1993). Lymphotxin β , a novel member of the TNF family that forms a heteromeric complex with lymphotxin on the cell surface. *Cell* 72, 847-858.
- Chomczynski, P., and Sacchi, N. (1987). Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal. Biochem.* 162, 156-159.
- Cohen, J. J., Duke, R. C., Fadock, V. A., and Selina, K. S. (1992). Apoptosis and programmed cell death in immunity. *Annu. Rev. Immunol.* 10, 267-293.
- Cohen, P. L., and Elsenberg, R. A. (1991). *Lpr* and *gld*: single gene models of systemic autoimmunity and lymphoproliferative disease. *Annu. Rev. Immunol.* 9, 243-269.
- Compton, M. M. (1992). A biochemical hallmark of apoptosis: internucleosomal degradation of the genome. *Cancer Metastasis Rev.* 11, 105-110.
- Conzelmann, A., Cortés, P., Cianfriga, M., Silva, A., and Nabholz, M. (1982). Hybrids between rat lymphoma and mouse T cells with inducible cytolytic activity. *Nature* 298, 170-172.
- Dhein, J., Daniel, P. T., Trauth, B. C., Oehm, A., Möller, P., and Krammer, P. H. (1992). Induction of apoptosis by monoclonal antibody anti-APO-1 class switch variants is dependent on cross-linking of APO-1 cell surface antigens. *J. Immunol.* 149, 3166-3173.
- Dower, W. J., Miller, J. F., and Pascale, C. W. (1988). High efficiency transformation of E. coli by high voltage electroporation. *Nucl. Acids Res.* 16, 6127-6145.
- Eck, M. J., and Sprang, S. R. (1989). The structure of tumor necrosis factor- α at 2.8 Å resolution. *J. Biol. Chem.* 264, 17595-17605.
- Eck, M. J., Utsch, M., Rinderknecht, E., de Vos, A. M., and Sprang, S. R. (1992). The structure of human lymphotxin (tumor necrosis factor- β) at 1.9-Å resolution. *J. Biol. Chem.* 267, 2119-2122.
- Ellis, R. E., Yuen, J., and Horvitz, H. R. (1991). Mechanisms and functions of cell death. *Annu. Rev. Cell Biol.* 7, 663-698.
- Fukunaga, R., Ishizaka-Kresia, E., Sato, Y., and Nagata, S. (1990). Expression cloning of a receptor for murine granulocyte colony-stimulating factor. *Cell* 61, 341-350.
- Golstein, P., Ojcius, D. M., and Young, J. D. E. (1991). Cell death mechanisms and the immune system. *Immunol. Rev.* 121, 29-65.
- Goodwin, R. G., Alderson, M. R., Smith, C. A., Armitage, R. J., VandenBos, T., Jerzy, R., Tough, T. W., Schoenborn, M. A., Davis-Smith, T., Hennen, K., Falk, B., Cosman, D., Baker, E., Sutherland, G. R., Grabstein, K. H., Farnish, T., Gid, J. G., and Beckmann, M. P. (1993). Molecular and biological characterization of a ligand for CD27 defines a new family of cytokines with homology to tumor necrosis factor. *Cell* 73, 447-458.
- Harada, N., Castle, B. E., Gorman, D. M., Roh, N., Schueurs, J., Barrett, R. L., Howard, M., and Miyajima, A. (1990). Expression cloning of a cDNA encoding the murine interleukin 4 receptor based on ligand binding. *Proc. Natl. Acad. Sci. USA* 87, 857-861.
- Itoh, N., Yonehara, S., Ishii, A., Yonehara, M., Mizushima, S.-I., Semeshima, M., Hase, A., Seto, Y., and Nagata, S. (1991). The polypeptide encoded by the cDNA for human cell surface antigen Fas can mediate apoptosis. *Cell* 66, 233-243.
- Roh, N., Tsujimoto, Y., and Nagata, S. (1993). Effect of bcl-2 on Fas antigen-mediated cell death. *J. Immunol.* 151, 621-627.
- Jones, E. Y., Stuart, D. I., and Walker, N. P. C. (1989). Structure of tumor necrosis factor. *Nature* 338, 225-228.
- Ju, S.-T. (1991). Distinct pathways of CD4 and CD8 cells induce rapid target DNA fragmentation. *J. Immunol.* 146, 812-818.
- Kozak, M. (1991). An analysis of vertebrate mRNA sequences: initiation of translational control. *J. Cell Biol.* 115, 887-903.
- Kriegler, M., Perez, C., DeFay, K., Albert, I., and Lu, S. D. (1988). A novel form of human TNF/Cachectin is a cell surface cytotoxic transmembrane protein: ramifications for the complex physiology of TNF. *Cell* 53, 45-53.
- Leid, P. W., Zijdenfeld, A., Linders, K., Rudnicki, M. A., Janssch, R., and Berns, A. (1991). Simplified mammalian DNA isolation procedure. *Nucl. Acids Res.* 19, 4293.
- Loetscher, H., Gentz, R., Zulauf, M., Lustig, A., Tabuchi, H., Schlaeger, E.-J., Brockhaus, M., Gallas, H., Manneberg, M., and Lesslauer, W. (1991). Recombinant 55-kDa tumor necrosis factor (TNF) receptor. *J. Biol. Chem.* 266, 18324-18329.
- Meier, T., Arni, S., Matarikannan, S., Poincelot, M., and Hoessli, D. (1992). Immunodetection of biotinylated lymphocyte-surface proteins by enhanced chemiluminescence. A nonradioactive method for cell-surface protein analysis. *Anal. Biochem.* 204, 220-226.
- Misling, A. (1992). Germ-cell death during spermatogenesis in the testis of the golden hamster. *Cell Tissue Res.* 267, 583-590.
- Nagata, S. (1993). Apoptosis-mediated Fas antigen and its natural mutation. In *Apoptosis* B. L. D. Tomel and F. C. Cope, eds. (Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press), in press.
- Nishimura, Y., Yokoyama, M., Araki, K., Ueda, R., Kudo, A., and Watanabe, T. (1987). Recombinant human-mouse chimeric monoclonal antibody specific for common acute lymphocytic leukemia antigen. *Cancer Res.* 47, 999-1005.
- Oehm, A., Behrmann, I., Falk, W., Pawlis, M., Maier, G., Kies, C., Li-Weber, M., Richards, S., Dhein, J., Trauth, B. C., Ponzsingl, H., and Krammer, P. H. (1992). Purification and molecular cloning of the APO-1 cell surface antigen, a member of the tumor necrosis factor/tumor growth factor receptor superfamily: sequence identity with the Fas antigen. *J. Biol. Chem.* 267, 10709-10715.
- Ogasawara, J., Watanabe-Fukunaga, R., Adachi, M., Matsuzawa, A., Kasugai, T., Kikuchi, Y., Itoh, N., Suda, T., and Nagata, S. (1993). Lethal effect of the anti-Fas antibody in mice. *Nature* 364, 808-809.
- Old, L. J. (1987). Another chapter in the long history of endotoxin. *Nature* 330, 602-603.
- Ozdemirli, M., El-Khalil, M., Bastiani, L., Akdeniz, H., Kuchroo, V., and Ju, S.-T. (1992). The cytotoxic process of CD4 Th1 clones. *J. Immunol.* 148, 1889-1895.
- Pennica, D., Nedwin, G. E., Haylick, J. S., Seeburg, P. H., Derynck, R., Palladino, M. A., Kohr, W. J., Aggeler, B. B., and Goeddel, D. V. (1984). Human tumor necrosis factor: precursor structure, expression and homology to lymphotxin. *Nature* 312, 724-728.
- Perez, C., Albert, I., DeFay, K., Zachariades, N., Gooding, L., and Kriegler, M. (1990). A nonsecretable cell surface mutant of tumor necrosis factor (TNF) kills by cell-to-cell contact. *Cell* 63, 251-258.
- Podack, K. R., Hengartner, H., and Uchtenheld, M. G. (1991). A central role of perforin in cytotoxicity? *Annu. Rev. Immunol.* 9, 129-157.
- Potter, H., Web, L., and Leder, P. (1984). Enhancer dependent expression of human κ immunoglobulin genes introduced into mouse pre-B lymphocytes by electroporation. *Proc. Natl. Acad. Sci. USA* 81, 7161-7165.
- Raff, M. C. (1992). Social controls on cell survival and cell death. *Nature* 356, 397-400.
- Ramsdell, F., and Fowlkes, B. J. (1990). Clonal deletion versus clonal energy: the role of the thymus in inducing self tolerance. *Science* 248, 1342-1348.

- Rouvier, E., Luciani, M.-F., and Golstein, P. (1993). Fas involvement in Ca^{2+} -independent T cell-mediated cytotoxicity. *J. Exp. Med.* 177, 195-200.
- Russell, J. H., Rush, B., Weaver, C., and Wang, R. (1993). Mature T cells of autoimmune *lpr/lpr* mice have a defect in antigen-stimulated suicide. *Proc. Natl. Acad. Sci. USA* 90, 4409-4413.
- Sambrook, J., Fritsch, E. F., and Maniatis, T. (1989). *Molecular Cloning: A Laboratory Manual*, Second Edition (Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press).
- Sedivy, J. M. (1988). New genetic methods for mammalian cells. *Biol. Technology* 6, 1192-1196.
- Smith, R. A. and Baglioni, C. (1987). The active form of tumor necrosis factor is a trimer. *J. Biol. Chem.* 262, 6951-6954.
- Smith, C. A., Grusa, H.-J., Davis, T., Anderson, D., Farrah, T., Baker, E., Sutherland, G. R., Brannan, C. I., Copeland, N. G., Jenkins, N. A., Grabstein, K. H., Gliniak, B., McAlister, I. B., Fanslow, W., Alderson, M., Falk, B., Gimpel, S., Gills, S., Din, W. S., Goodwin, R. G., and Arnhege, R. J. (1993). CD30 antigen, a marker for Hodgkin's lymphoma, is a receptor whose ligand defines an emerging family of cytokines with homology to TNF. *Cell* 73, 1349-1360.
- Trauth, B. C., Kna, C., Peters, A. M. J., Matzku, S., Möller, P., Falk, W., Debatin, K.-M., and Krammer, P. H. (1989). Monoclonal antibody-mediated tumor regression by induction of apoptosis. *Science* 245, 301-305.
- Uetsuki, T., Naito, A., Nagata, S., and Kazuo, Y. (1989). Isolation and characterization of the human chromosomal gene for polypeptide chain elongation factor 122a. *J. Biol. Chem.* 264, 5791-5798.
- Walker, N. I., Harmon, B. V., Gobe, G. C., and Kerr, J. F. R. (1988). Patterns of cell death. *Methods Achiev. Exp. Pathol.* 13, 18-54.
- Watanabe-Fukunaga, R., Brannan, C. I., Copeland, N. G., Jenkins, N. A., and Nagata, S. (1992a). Lymphoproliferation disorder in mice explained by defects in Fas antigen that mediates apoptosis. *Nature* 356, 314-317.
- Watanabe-Fukunaga, R., Brannan, C. I., Itoh, N., Yonehara, S., Copeland, N. G., Jenkins, N. A., and Nagata, S. (1992b). The cDNA structure, expression, and chromosomal assignment of the mouse Fas antigen. *J. Immunol.* 148, 1274-1279.
- Wyllie, A. H., Kerr, J. F. R., and Currie, A. R. (1980). Cell death: the significance of apoptosis. *Int. Rev. Cytol.* 68, 251-308.
- Yonehara, S., Ishii, A., and Yonehara, M. (1989). A cell-killing monoclonal antibody (anti-Fas) to a cell surface antigen co-downregulated with the receptor of tumor necrosis factor. *J. Exp. Med.* 169, 1747-1756.
- Zhou, T., Bluethmann, H., Eldridge, J., Berry, K., and Mountz, J. D. (1993). Origin of CD4⁺CD8⁺B220⁺ T cells in MRL-*lpr/lpr* mice. *J. Immunol.* 150, 3651-3657.

GenBank Accession Number

The accession number for the cDNA sequence reported in this paper is U03470.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.